

Large N Gauge Theories and Anti-de Sitter Bag Model

I.V. Volovich *

*Centro Vito Volterra
Universita di Roma Tor Vergata, Rome, Italy*

Abstract

Using a proposal of Maldacena one describes the large N limit of gauge theories in terms of supergravity solutions on anti-de Sitter space. From this point of view we discuss a possible scenario for quark confinement in gauge theory by describing hadrons as strongly curved universes. In particular an interpretation of black hole as a bag model in SQCD is discussed. One relates the mystery of curvature singularities in classical general relativity with the mystery of quark confinement. The AdS bag model is defined by computing the probe membrane action in supergravity background. It naturally implies the "Cheshire Cat bag" principle. The confining pressure in the MIT bag model is related with the cosmological constant in the AdS bag model. The Skyrme model is interpreted as an effective theory describing black holes.

*Permanent address: Steklov Mathematical Institute, Gubkin St.8, GSP-1, 117966, Moscow, Russia, volovich@mi.ras.ru

In many recent works a relation between the large N limit of supersymmetric Yang-Mills theories and supergravity has been explored [1]-[42]. It was proposed by Maldacena [1] that in particular the large N limit of the $\mathcal{N} = 4$ super Yang-Mills theory in four dimensions with $U(N)$ gauge group is equivalent to the type IIB string theory on $AdS_5 \times S^5$. The gauge theory describes a region of spacetime near the horizon of a black hole which is known to have anti-de Sitter (AdS) geometry. Moreover Horowitz and Ooguri [15] have argued that the gauge theory actually describes string theory on both sides of the horizon. Therefore one has to find in gauge theory an analogue of the horizon which is the basic notion of the black hole physics.

In this note we discuss a scenario for a very natural possible correspondence between basic notions of gauge theory and gravity. It is tempting to speculate that confinement in gauge theory corresponds to black holes in gravity and one can try to describe hadrons as strongly curved universes or black holes. In this picture the horizon corresponds to the surface of the bag in a bag model. One relates the mystery of curvature singularities in classical general relativity with the mystery of quark confinement.

It is proposed in [34] that the gauge theory could provide a resolution of the singularity inside the black hole since it is perfectly regular. From our point of view the gauge theory is not a regular theory, rather it is a very hard theory especially in the infra-red regime. One believes that perhaps the gauge theory could really give some help for understanding the curvature singularities in general relativity but only if we are able to understand the problem of confinement.

The bag model was one of the first attempts to understand the color confinement [43, 44]. In bag models quarks and gluons are confined to a bounded region that usually is taken as a static sphere of radius R . In the MIT bag model [44] the bag exterior acts as a perfect color dielectric, while the color degrees of freedom are confined to the bag interior. At the surface of the bag, the gluons are subject to the confining boundary conditions

$$nE^a = 0, \quad n \times B^a = 0 \quad (1)$$

where E^a and B^a are color electric and magnetic fields and n is the outward normal to the bag surface. For the quarks, the confining boundary condition states that there is no flux of the isoscalar current through the bag surface:

$$n_i \bar{q} \gamma^i q = 0 \quad (2)$$

The bag model gives a rather satisfactory description of masses, magnetic momenta and some other properties of hadrons. One of the main problems with the bag model is that it is unclear how to derive the basic assumptions of the model from QCD.

In the "Cheshire Cat bag" approach [45] the bag itself has no physical significance. One just formally separates the space-time in a bag region where QCD still acts and an exterior region where QCD is integrated out by an effective action formulated in hadronic degrees of freedom (for example the Skyrme model), and a boundary. The location of the boundary has no physical consequences and moreover fermionic degrees of freedom can be replaced by bosonic ones.

The idea of describing elementary particles as strongly curved universes was discussed by Frolov and Markov [46], by Isham, Salam and Strathdee [47] and other

authors, see for example [48]. Black holes and negatively curved anti-de Sitter space could evidently provide baglike structures. Timelike geodesics in AdS are known periodic with frequency $1/R$ where R is the AdS radius. Therefore if one assumes that free quarks move along geodesics then one gets that they carry out harmonic oscillations within a sphere of radius R .

According to [1] the large N limit of supersymmetric gauge theory is described by supergravity on AdS space. Similar description was also proposed for large N gauge theories without supersymmetry [16] including QCD. It is known that QCD in the large N limit is equivalent to a meson theory which is not reduced to the non-linear sigma model or Skyrme model. It is a complex unknown theory with an infinite number of elementary fields [50]. Now by using the Maldacena conjecture one can try to describe this unknown meson theory in term of the infinite number of Kaluza-Klein modes. Probably in this way one gets the "Kaluza-Klein corrections" to the sigma model and to the Skyrme model. A gauge invariant version of the Skyrme model is considered in [51].

In the hybrid chiral bag model [52] a surface coupling with an external pion field is introduced to restore chiral invariance which was broken in the original MIT bag model. The Lagrangian for the external pion field is that of the non-linear σ -model or the Skyrme model. A specific solution was used for the meson field where its isospin direction point radially, the so called hedgehog solution. The same solution arises in a model for baryons, originally proposed by Skyrme and then revisited by Witten [50]. In this model the baryons are topologically stable solutions in a non-linear σ -model describing only meson fields. The topological charge of the soliton solution is identified with the baryon number.

In the recent papers [16, 21, 22, 38, 39, 40] the thermodynamics of black holes in AdS space was used for the investigation of thermodynamics of gauge theories. An ansatz for the Wilson loop in the large N limit was proposed which involves calculating the area of a probe string worldsheet in a supergravity background.

The ansatz for the Wilson loop

$$W(C) = \frac{1}{N} \langle Tr P e^{\int_C A} \rangle \quad (3)$$

used in [21, 22] is

$$W(C) = e^{-I} \quad (4)$$

where

$$I = \int d^2\sigma \sqrt{\det h_{ab}}, \quad h_{ab} = g_{MN} \partial_a x^M \partial_b x^N \quad (5)$$

The action I is the area of the string worldsheet which at the boundary of AdS describes the loop C . The metric g_{MN} is a Schwarzschild-de-Sitter supergravity solution describing the near horizon geometry of near extremal D3-branes in type IIB string theory

$$ds^2 = \alpha' \left[\frac{U^2}{R^2} (-f dt^2 + dx_i^2) + \frac{R^2}{f U^2} dU^2 + R^2 d\Omega_5^2 \right] \quad (6)$$

where $f = 1 - U_0^4/U^4$, $R^2 = g_{YM}^2 N$ and U_0^4 is proportional to the energy density on the brane.

From this it seems natural to propose that the computation of the volume of a probe membrane worldvolume in a supergravity background gives an *AdS* bag model. It naturally allows the "Cheshire Cat" principle [45]. A proposal for the expectation value of the bag operator $B(M)$ [49] is

$$\langle B(M) \rangle = e^{-S}. \quad (7)$$

Here M is a two-dimensional surface describing the boundary of the bag (membrane) and S is the volume of the membrane which at the boundary of *AdS* describes the surface M ,

$$S = \int d^3\sigma \sqrt{\det h_{ab}}, \quad h_{ab} = g_{MN} \partial_a x^M \partial_b x^N \quad (8)$$

In this picture mesons in QCD correspond to black holes in gravity and the σ -model or the Skyrme model should give an effective description of black holes because they give a phenomenological description of mesons. A gravitational analogue of barions in such an approach is not clear at the moment. The boundary conditions (1) and (2) are set on the spacelike surface. The horizon of a black hole is a light-like surface. Hence an analogue of (1) and (2) one has to set on a stretched horizon.

A derivation of the phenomenological description of multiple black holes one can get along the discussion in [26] where bunches of branes are considered. The supergravity solution in type IIB string theory carrying D3-brane charge has the form

$$ds^2 = f^{-1/2}(-dt^2 + dx_i^2) + f^{1/2}dy_\mu^2 \quad (9)$$

where $i = 1, 2, 3$, $\mu = 4, 5, 6, 7, 8, 9$ and $f = f(y)$ is a harmonic function.

If we consider two bunches of D3-branes located at points $y^{(1)}$ and $y^{(2)}$ then

$$f = 1 + 4\pi g\alpha' \left[\frac{N_1}{|y - y^{(1)}|^4} + \frac{N_2}{|y - y^{(2)}|^4} \right] \quad (10)$$

This metric describes two black holes. There are gravitational forces between two bunches and it seems natural to think that this configuration of branes is described by super Yang-Mills theory in the curved background. From the previous discussion one suggests that these black holes can be considered as mesons and can be described by using the sigma model or the Skyrme model lagrangians. The same reasoning can be applied to a configuration of an arbitrary number of non-extremal black holes. The quantum Boltzmann statistics found in hadron physics is relevant also to the physics of black holes, see [6, 26] and refs therein. In the case $gN_1 \gg gN_2 \gg 1$ one has super Yang-Mills theory in the 3-brane background.

I am grateful to I. Aref'eva for useful comments. The work is supported in part by RFFI grant 96-01-00312 and INTAS grant 96-0698.

References

- [1] J. Maldacena, hep-th/9711200
- [2] S. Ferrara and C. Fronsdal, hep-th/9712239; hep-th/9802126.
- [3] S. S. Gubser, I. R. Klebanov and A. W. Peet, Phys. Rev. **D54**(1996) 3915.
- [4] I. R. Klebanov, Nucl. Phys. **B496** (1997) 231.
- [5] S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, hep-th/9703040
- [6] I.V. Volovich, hep-th/9608137; hep-th/9705013.
- [7] K. Sfetsos and K. Skenderis, hep-th/9711138,
- [8] R. Kallosh, J. Kumar and A. Rajaraman, hep-th/9712073.
- [9] H.J.Boonstra, B.Peeters and K. Skenderis, hep-th/9801076.
- [10] P. Claus, R. Kallosh, J. Kumar, P. Townsend, and A. van Proeyen, hep-th/9801206.
- [11] N. Itzhaki, J. M. Maldacena, J. Sonnenschein, and S. Yankielowicz, hep-th/9802042.
- [12] M. Gunaydin and D. Minic, hep-th/9802047.
- [13] I.Ya. Aref'eva and I.V. Volovich, hep-th/9802091
- [14] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, hep-th/9802109.
- [15] G. T. Horowitz and H. Ooguri, hep-th/9802116.
- [16] E. Witten, hep-th/9802150
- [17] S. Kachru and E. Silverstein, hep-th/9802183
- [18] M. Berkooz, hep-th/9802195.
- [19] V. Balasubramanian and F. Larsen, hep-th/9802198
- [20] S. Ferrara, C. Fronsdal and A. Zaffaroni, hep-th/9802203.
- [21] S.-J. Rey and J. Yee, hep-th/9803001
- [22] J. Maldacena, hep-th/9803002
- [23] M. Flato and C. Fronsdal, hep-th/9803013
- [24] A. Lawrence, N. Nekrasov and C. Vafa, hep-th/9803015
- [25] S.S. Gubser, A. Hashimoto, I.R. Klebanov and M. Krasnitz, hep-th/9803023.
- [26] I. Ya. Aref'eva and I. V. Volovich, hep-th/9803028.
- [27] L. Castellani, A. Ceresole, R. D'Auria, S. Ferrara, P. Fré and M. Trigiante, hep-th/9803039.

- [28] O. Aharony, Y. Oz and Z. Yin, hep-th/9803051.
- [29] S. Minwalla, hep-th/9803053.
- [30] M.J.Duff, H.Lu, C.N.Pope, hep-th/9803061.
- [31] R. G. Leigh and M. Rozali, hep-th/9803068.
- [32] M. Bershadsky, Z. Kakushadze, and C. Vafa, hep-th/9803076.
- [33] A. Rajaraman, hep-th/9803082.
- [34] G.T. Horowitz and S.F. Ross, hep-th/9803085.
- [35] S. Ferrara, A. Kehagias, H. Partouche and A. Zaffaroni, hep-th/9803109.
- [36] J. A. Minahan, hep-th/9803111.
- [37] J. Gomis, hep-th/9803119.
- [38] E. Witten, hep-th/9803131
- [39] S.J. Rey, S. Theisen and J.T. Yee, hep-th/9803135
- [40] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, hep-th/9803137
- [41] M. Gunaydin, hep-th/9803138
- [42] Y.Oz and J.Terning, hep-th/9803167
- [43] P.N. Bogoliubov, Ann. Inst. Henri Poincare, 8(1967)163
- [44] A. Chodos, R.L. Jaffe, K. Johnson, K.B. Thorn and V.F. Weisskopf, Phys. Rev. D9(1974)3471
- [45] S. Nadkarni, H.B. Nielsen and I.Zahed, Nucl. Phys. B253(1985)308; H.B. Nielsen, M. Rho, A. Wirzba and I.Zahed, Phys. Lett. B281(1992)345
- [46] V.P. Frolov and M.A. Markov, Theor. Math. Phys. 17(1970)249
- [47] C. Isham, A. Salam and J. Strathdee, Phys. Rev. D3(1971)867; A. Salam and J. Strathdee, Phys. Rev. D18(1978)4596
- [48] C.Sivaram and K.P.Sinha, Phys. Rep. 51(1979)111 ; Yu.A.Golfand, JETP,43(1962)256; N.A.Chernikov and E.A.Tagirov, Ann. Inst. H.Poincare,9(1968)109; L.Castell, Nuvo Cim.61A(1968)585; E.van Beveren, C.Dullemond and T.A.Rijken, Phys. Rev. D30(1984)103; H.C.Rosu, Nuovo Cim. B107(1993)313.
- [49] I.Ya. Aref'eva, Phys. Lett. 92B(1980)51
- [50] E. Witten, Nucl.Phys. B160 (1979) 57; B223 (1983) 433.
- [51] L.D. Faddeev, Lett. Math. Phys. 1(1976)289
- [52] A. Chodos and C.B. Thorn, Phys.Rev. D12 (1975) 2733